ACR REVIEW 2026

#ACReview26

Lo mejor de ACR



Sociedad Española de Reumatología

PATROCINADO POR: AstraZeneca · Gebro Pharma · Lilly A MEDICINE COMPANY

ACR REVIEW 2026

Lo mejor de LES, Sjögren y Esclerosis Sistémica

Cristiana Sieiro Santos

Centre for Musculoskeletal Research, School of Biological Sciences,
Faculty of Biology, Medicine and Health,
University of Manchester, Manchester, UK


Manchester University NHS Foundation Trust (MFT)

Sociedad Española de Reumatología

PATROCINADO POR: AstraZeneca Gebro Pharma Lilly A MEDICINE COMPANY

ABSTRACT NUMBER: 1222


Histological Remission and Response in Severe and Refractory Lupus Nephritis Patients after Treatment with Rese-cel (Resecabtagene Autoleucel), an AutologousCD19-CAR T Cell Therapy



ABSTRACT NUMBER: 1862

Post-Treatment Proteinuria Imperfectly Predicts Histological Remission in Lupus Nephritis: Repeat Biopsy Data from ReBioLup Support Reappraisal of Trial Endpoints

Diseño del Estudio y Población



- Medianas de edad
- Duración del LES
- Clases histológicas iniciales (III, IV, III/IV+V, V)

Resultados Clave: Discordancia Clínica




Proteinuria <0,5 g/g → Actividad histológica significativa (AI≥4): **17%**

Proteinuria >1 g/g → Actividad histológica mínima o ausente (AI<3): **10,6%**

Pacientes con AI≥4 cumplían criterios habituales de respuesta renal completa: **66,7%**

Gráficos ROC: Predicción de Remisión Histológica



UPCR = 0,22 g/g para AI=0 (AUC=0,79)

UPCR = 0,25 g/g para AI≤1 (AUC=0,84)


La proteinuria a los 12 meses predice la remisión histológica sólo parcialmente. Puntos de corte óptimos identificados, pero con traslape y discordancia clínica relevante.

Mensaje Principal (Conclusión)

Existe una discordancia clínicamente relevante entre la respuesta renal basada en proteinuria y la remisión histológica. Proteinuria baja no garantiza remisión, y proteinuria persistente no implica inflamación activa. Se deben reconsiderar los umbrales de respuesta renal en ensayos clínicos de nefritis lúpica.


ABSTRACT NUMBER: 1857

Progression of Systemic Activity, Organ Damage, and Symptom Burden in Sjögren's Disease: Results from the Prospective Multicenter SjögrenSER-PROS Registry




ABSTRACT NUMBER: 0986

Induced pluripotent stem cell (iPSC)-derived cardiac microtissues as a modeling and drug testing platform for systemic sclerosis primary heart involvement




ABSTRACT NUMBER: 1934

Proteome-wide autoantibody profiling defines serology-based subtypes of gastrointestinal disease in systemic sclerosis




ABSTRACT NUMBER: 1189

Gut Microbiome and Metabolomic Signatures Of Progressive Pulmonary Fibrosis in Systemic Sclerosis-Associated Interstitial Lung Disease: Insights from an International Cohort



ABSTRACT NUMBER: 0981

Efficacy and Safety of Sotatercept in Patients with Pulmonary Arterial Hypertension (PAH) Associated with CTD: A Pooled Analysis of Three Phase 3 Trials



ACR REVIEW 2026

#ACReview26

Lo mejor de ACR



Sociedad Española de Reumatología

PATROCINADO POR: AstraZeneca · Gebro Pharma · Lilly A MEDICINE COMPANY